



6<sup>th</sup> IAS CONFERENCE  
ON HIV PATHOGENESIS,  
TREATMENT AND PREVENTION

17-20 JULY 2011 – ROME, ITALY



## Abstract A-361-0084-02626

Decrease of Ki-67 proliferation marker correlates with CD4+ cell count increase and predicts viral load suppression following 4 week AV-HALT treatment: clinical Proof-of-Concept of an immunological predictive marker.

D. De Forni<sup>1</sup>, E. Katabira<sup>2</sup>, R. Maserati<sup>3</sup>, P. Cahn<sup>4</sup>, D. V. Baev<sup>1</sup>, B. Poddesu<sup>1</sup>, M.R. Stevens<sup>5</sup>, F.Lori<sup>1</sup>

<sup>1</sup> ViroStatics srl, Alghero, Italy, <sup>2</sup> Makerere Medical School, Kampala, Uganda, <sup>3</sup> Fondazione IRCCS Policlinico San Matteo, Infectious Diseases Unit, Pavia, Italy, <sup>4</sup> Fundación Huesped, Buenos Aires, Argentina, <sup>5</sup> ViroStatics srl, Princeton, United States

**Background:** Chronically excessive immune system activation plays a central role in HIV-AIDS pathogenesis. Although HAART can maximally suppress HIV viral load, T cell activation and proliferation markers do not return to non-HIV-infected values. An immunological biomarker predictive of viral suppression could be of value to clinicians selecting antiretroviral regimens. We tested whether pre-therapy Ki-67 proliferation marker values in naïve and Central Memory (CM) CD4+ T cells could predict HIV suppression by a newly-developed class of anti-HIV drugs known as AntiViral-HyperActivation Limiting Therapeutics (AV-HALTs) exhibiting dual activities: suppressing HIV replication and limiting excessive T cell proliferation.

**Methods:** Blood samples from 32 subjects receiving the AV-HALT VS411, a novel combination of an antiviral (low-dose, slow-release 2',3'-dideoxyinosine) and an antiproliferative drug (low-dose hydroxycarbamide), were analyzed in a multinational 4-week Phase IIa, dose-finding study using 10-color flow cytometry including T cell subsetting, activation and proliferation markers. Pearson correlation was employed for statistical analysis.

**Results:** At Day 28, the median decrease in HIV RNA was 1.47 log<sub>10</sub> with only two of the 32 subjects reaching < 50 copies/mL; median CD4+ T cell increase was +108 cells/mm<sup>3</sup>. Significant correlations were found between the percentage of Ki-67+ naïve (CD45RA+CD27+) CD4+ T cells and CD4+ T cell counts at baseline and following treatment with  $r^2=0.191$  and  $r^2=0.259$ , respectively ( $P < .05$ ). A predictive correlation ( $r^2= 0.262$ ,  $P < .05$ ) between the percentage of Ki-67+ CM (CD45RA-CD27+) CD4+ T cells at baseline and log<sub>10</sub> HIV viral load changes observed after 28 days of AV-HALT therapy was also found.

**Conclusion:** Pre-treatment expression of the proliferation marker Ki-67 in Central Memory CD4+ T cells was predictive of viral load decline over 28 days of AV-HALT therapy. The study provides the clinical Proof-of-Concept that percentage of Ki-67+ Central Memory CD4+ T cells at baseline is predictive of HIV suppression following AV-HALT treatment.